

FILTERED HOPF ALGEBRAS AND COUNTING GEODESIC CHORDS

URS FRAUENFELDER AND FELIX SCHLENK

ABSTRACT. We prove lower bounds on the growth of certain filtered Hopf algebras by means of a Poincaré–Birkhoff–Witt type theorem for ordered products of primitive elements.

When applied to the loop space homology algebra endowed with a natural length-filtration, these bounds lead to lower bounds for the number of geodesic paths between two points. Specifically, given a closed manifold M whose universal covering space is not homotopy equivalent to a finite complex and whose fundamental group has polynomial growth,

for any Riemannian metric on M , any pair of non-conjugate points $p, q \in M$, and every component \mathcal{C} of the space of paths from p to q , the number of geodesics in \mathcal{C} of length at most T grows at least like $e^{\sqrt{T}}$.

Using Floer homology, we extend this lower bound to Reeb chords on the spherisation of M , and give a lower bound for the volume growth of the Reeb flow.

1. INTRODUCTION AND MAIN RESULTS

Consider a closed connected manifold M . Then M is said to be of *finite type* if its universal cover \widetilde{M} is homotopy equivalent to a finite CW-complex. Equivalently, M is *not* of finite type if there exists $k \in \{2, \dots, \dim M\}$ such that $H_k(\widetilde{M})$ is not finitely generated. Let $\mathfrak{m} = \mathfrak{m}(M)$ be the minimal such k .

A finitely generated group G is called *polycyclic* if it admits a subnormal series with cyclic factors. Moreover, G is *virtually polycyclic* if it has a polycyclic subgroup of finite index.

Choose a point $p \in M$ and let $\Omega_0 M$ be the space of contractible continuous loops in M based at p . For $T > 0$ denote by $\Omega_0^T M$ the space of contractible piecewise smooth loops based at p whose length (with respect to a fixed Riemannian metric on M) is at most T . The inclusions $\iota^T: \Omega_0^T M \hookrightarrow \Omega_0 M$ induce the maps $\iota_*^T: H_*(\Omega_0^T M) \rightarrow H_*(\Omega_0 M)$ in homology. Note that the image $\iota_*^T H_*(\Omega_0^T M)$ is the part of the homology of $\Omega_0 M$ that is generated by cycles made of piecewise smooth loops of length $\leq T$.

We say that a real-valued function f defined on \mathbb{N} or on $\mathbb{R}_{>0}$ *grows at least like* $e^{\sqrt{T}}$ if there exists a constant $c > 0$ such that $f(T) \geq ce^{\sqrt{T}}$ for all large enough T .

Date: June 18, 2014.

UF partially supported by the Basic Research fund 2013004879 of the Korean government.

FS partially supported by SNF grant 200020-144432/1.

2000 *Mathematics Subject Classification.* Primary 16T05, Secondary 37B40, 37C35, 53D25, 57T25.

Theorem 1.1. *Let M be a closed connected manifold that is not of finite type and has virtually polycyclic fundamental group. Assume also that there exists a field \mathbb{F} such that $H_{\mathfrak{m}}(\widetilde{M}; \mathbb{F})$ is infinite-dimensional. Then the function of T*

$$\dim \iota_*^T H_*(\Omega_0^T M; \mathbb{F})$$

grows at least like $e^{\sqrt{T}}$.

Remarks 1.2. 1. The assumption that $H_{\mathfrak{m}}(\widetilde{M}; \mathbb{F})$ is infinite-dimensional is equivalent to the assumption that $\pi_{\mathfrak{m}}(M) \otimes \mathbb{F}$ is infinite-dimensional (see Section 5.4 for the proof).

2. During the proof we shall see that $H_{\mathfrak{m}-1}(\Omega_0 M; \mathbb{F})$ is infinite-dimensional and that

$$\dim H_{(\mathfrak{m}-1)n}(\Omega_0 M; \mathbb{F}) \geq q(n) \quad \text{for } n \geq 2,$$

where $q(n)$ is the number of partitions of n into distinct parts. The fact that this function grows like $e^{\sqrt{n}}$ explains the lower bound in Theorem 1.1.

3. Our assumption that there exists a field \mathbb{F} such that $H_{\mathfrak{m}}(\widetilde{M}; \mathbb{F})$ is infinite-dimensional not only plays an important role in our proof, but may well be essential. Indeed, in Section 6 we first generalize Theorem 1.1 to compact simplicial complexes embedded in some Euclidean space, and then show that a simplicial complex K homotopy equivalent to the mapping torus of a degree- p map of S^n (with p prime and $n \geq 2$) has the following properties: All assumptions of Theorem 1.1 are met besides for $H_{\mathfrak{m}}(\widetilde{K}; \mathbb{F})$ being zero- or one-dimensional for every field \mathbb{F} , but $\iota_*^T H_*(\Omega_0^T K)$ is a free Abelian group whose rank grows not more than linearly with T .

For *manifolds* of non-finite type it is still possible that the rank of $\iota_*^T H_*(\Omega_0^T M)$ grows exponentially (cf. the Question in [27, p. 289] and the discussion in [8, §7.1]).

Examples 1.3. There are many manifolds meeting the assumptions of Theorem 1.1. For instance, take closed connected manifolds M_1 and M_2 of the same dimension, with $\pi_1(M_1)$ an infinite polycyclic group and M_2 simply-connected and not a homotopy sphere. For simplicity we also assume that $H_2(M_2) \neq 0$. Consider the connected sum $M = M_1 \# M_2$. Then $\pi_1(M) = \pi_1(M_1)$ is polycyclic. The universal cover \widetilde{M} is the universal cover \widetilde{M}_1 with one copy of M_2 attached for each element of $\pi_1(M_1)$. Since $\pi_1(M_1)$ is infinite, M is not of finite type, and $\mathfrak{m}(M) = 2$. Choose a field \mathbb{F} such that $H_{\mathfrak{m}}(M_2; \mathbb{F}) \neq 0$. Then $H_{\mathfrak{m}}(\widetilde{M}; \mathbb{F})$ is infinite-dimensional.

For any closed manifold homotopy equivalent to a connected sum $M = M_1 \# M_2$ with $\pi_1(M_1)$ of order at least three and with M_2 simply-connected and not a homotopy sphere, it has been shown in [26, 27] by a different method¹ that $\dim \iota_* H_*(\Omega_0^T M; \mathbb{F})$ grows even exponentially. It would therefore be interesting to find manifolds of non-finite type that are not homotopy equivalent to a non-trivial connected sum. There are obstructions to

¹Paternain and Petean consider a suitable finite simply-connected subcomplex $K \subset \widetilde{M}$ and prove exponential homological growth of the based loop space of K by showing exponential growth of the cohomology of the *free* loop space of K , which is isomorphic to the Hochschild homology of the singular cochain complex of K .

split a manifold (up to homotopy type) into a non-trivial connected sum, see [28, §7.4], but unfortunately these invariants are difficult to compute.

Question 1.4. *How large is the class of closed manifolds of non-finite type that are not homotopy equivalent to a non-trivial connected sum?*

1.1. Applications. Lower bounds for the rank of the homology of the sublevel sets $\Omega^T M$ are of interest because they classically lead, by Morse theory, to lower bounds for the number of geodesics of length $\leq T$ between non-conjugate points. Somewhat less classically, they also lead to lower bounds for the topological entropy of geodesic flows. Moreover, adding the tool of Floer homology, one gets lower bounds for the number of Reeb chords and for the topological entropy of Reeb flows on spherizations.

Before stating our two corollaries, we briefly recall what Reeb flows on spherizations are. Details can be found in the introduction of [8].

Reeb flows on spherizations. Consider a closed manifold M . The positive real numbers \mathbb{R}_+ freely act on the cotangent bundle T^*M by $r(q, p) = (q, rp)$. While the canonical 1-form $\lambda = pdq$ on T^*M does not descend to the quotient $S^*M := T^*M/\mathbb{R}_+$, its kernel does and defines a contact structure ξ on S^*M . We call the contact manifold (S^*M, ξ) the *spherization* of M . This contact manifold is co-orientable. The choice of a nowhere vanishing 1-form α on S^*M with $\ker \alpha = \xi$ (called a *contact form*) defines a vector field R_α (the *Reeb vector field* of α) by the two conditions $d\alpha(R_\alpha, \cdot) = 0$, $\alpha(R_\alpha) = 1$. Its flow φ_α^t is called the *Reeb flow* of α .

To give a more concrete description of the manifold (S^*M, ξ) and the flows φ_α^t , consider a smooth hypersurface Σ in T^*M which is *fiberwise starshaped* with respect to the zero-section: For every $q \in M$ the set $\Sigma_q := \Sigma \cap T_q^*M$ bounds a set in T_q^*M that is strictly starshaped with respect to the origin of T_q^*M . The hyperplane field $\xi_\Sigma := \ker(\lambda|_\Sigma)$ is a contact structure on Σ , and the contact manifolds (S^*M, ξ) and (Σ, ξ_Σ) are isomorphic.

Let φ_Σ^t be the Reeb flow on Σ defined by the contact form $\lambda|_\Sigma$. The set of Reeb flows on (S^*M, ξ) can be identified with the Reeb flows φ_Σ^t on the set of fiberwise starshaped hypersurfaces Σ in T^*M . The flows φ_Σ^t are restrictions of Hamiltonian flows: Consider a Hamiltonian function $H: T^*M \rightarrow \mathbb{R}$ such that $\Sigma = H^{-1}(1)$ is a regular energy surface and such that H is fiberwise homogeneous of degree one near Σ . For the Hamiltonian flow φ_H^t we then have $\varphi_H^t|_\Sigma = \varphi_\Sigma^t$. It follows that geodesic flows and Finsler flows (up to the time change $t \mapsto 2t$) are examples of Reeb flows on spherizations. Indeed, for geodesic flows the Σ_q are ellipsoids, and for (symmetric) Finsler flows the Σ_q are (symmetric and) convex. The flows φ_Σ^t for varying Σ are very different, in general, as is already clear from looking at geodesic flows on a sphere. In this paper we give uniform lower bounds for the growth of Reeb chords and for the complexity of all these flows on (S^*M, ξ) for manifolds M as in Theorem 1.1.

Growth of Reeb chords between two fibers. We say that $p, q \in M$ are *non-conjugate points* of the Reeb flow φ_α if $\bigcup_{t>0} \varphi_\alpha^t(S_p^*M)$ is transverse to S_q^*M . Given $p \in M$, the set of $q \in M$ that are non-conjugate to p has full measure in M by Sard's theorem. This notion

of being non-conjugate generalizes the one in Riemannian geometry (defined in terms of Jacobi fields).

For $p, q \in M$ denote by \mathcal{P}_{pq} the space of continuous paths in S^*M from S_p^*M to S_q^*M , and by $\Omega_{pq}M$ the space of continuous paths in M from p to q . We shall assume throughout that $\dim M \geq 3$, since otherwise M is of finite type. The fibers S_q^*M of the projection $pr: S^*M \rightarrow M$ are then simply connected, and so pr induces a bijection on the components of \mathcal{P}_{pq} and $\Omega_{pq}M$. The space $\Omega_{pq}M$ is homotopy equivalent to $\Omega_pM := \Omega_{p,p}M$, whose components are parametrized by the elements of the fundamental group $\pi_1(M, p)$.

Corollary 1.5. *Assume that M is not of finite type, has virtually polycyclic fundamental group, and that there is a field \mathbb{F} such that $H_{\mathfrak{m}}(\widetilde{M}; \mathbb{F})$ is infinite-dimensional. Let (S^*M, ξ) be the spherization of M . Then for any Reeb flow φ_α on (S^*M, ξ) , any pair of non-conjugate points $p, q \in M$ and every component \mathcal{C} of $\Omega_{pq}M$, the number of Reeb chords from S_p^*M to S_q^*M that belong to \mathcal{C} grows in time at least like $e^{\sqrt{T}}$.*

Remarks 1.6. 1. For the special case of geodesic flows, the time parameter equals the length run through. Hence the corollary translates to assertion (i) of the abstract.

To illustrate the corollary, we choose a Riemannian metric on M and a point $p \in M$. Let $C(p)$ be the cut locus of p . The subset $M \setminus C(p)$ is diffeomorphic to an open ball [23] and has full measure in M . For every $q \in M \setminus C(p)$ there is a unique shortest geodesic c_q from p to q . Call a path $\gamma \in \Omega_{pq}M$ *contractible* if $c_q^{-1} \circ \gamma$ is contractible in Ω_pM . The set U_p of points in $M \setminus C(p)$ that are not conjugate to p is also of full measure in M . Under the hypothesis of Corollary 1.5, for every $q \in U_p$ the number of contractible geodesics from p to q of length $\leq T$ grows at least like $e^{\sqrt{T}}$.

2. Virtually polycyclic groups are either virtually nilpotent or have exponential growth [34]. If the fundamental group $\pi_1(M)$ of a closed manifold M has exponential growth, then the number of Reeb chords from S_p^*M to S_q^*M grows exponentially in time for any, possibly conjugate, pair of points p, q (see [19, Corollary 1] for Reeb flows). Indeed, one finds one Reeb chord for each element of $\pi_1(M)$. In Corollary 1.5, however, we find “ $e^{\sqrt{T}}$ many” Reeb chords for each element of $\pi_1(M)$.

If one is only interested in the growth of Reeb chords from S_p^*M to S_q^*M , without specifying the component \mathcal{C} , then Corollary 1.5 is interesting only for virtually nilpotent fundamental groups, which by Gromov’s theorem from [12] are exactly those fundamental groups that grow polynomially. Indeed, it is believed that every finitely presented group that grows more than polynomially grows exponentially [9, Conjecture 11.3], and even for finitely generated groups of intermediate growth it is believed that they must grow at least like $e^{\sqrt{T}}$, cf. [10].

3. Let $\mu(\gamma)$ be the Conley–Zehnder index of a non-degenerate Reeb chord γ on $(S^*M; \alpha)$, normalized such that for geodesic flows $\mu(\gamma)$ is the Morse index of the non-degenerate geodesic γ (i.e. the number of conjugate points, counted with multiplicities, along γ). In the situation of Corollary 1.5, Remark 1.2.2 shows that for every component of $\Omega_{pq}M$ the number of Reeb chords from S_p^*M to S_q^*M of index $\mu(\gamma) = k$ is infinite if $k = \mathfrak{m} - 1$ and is at least the number $q(n)$ of partitions of n into distinct parts if $k = (\mathfrak{m} - 1)n$ and $n \geq 2$.

Intermediate volume growth. Consider a smooth diffeomorphism φ of a closed manifold X . Denote by \mathcal{S} the set of smooth compact submanifolds of X . Fix a Riemannian metric ρ on X , and denote by $\text{Vol}(\sigma)$ the j -dimensional volume of a j -dimensional submanifold $\sigma \in \mathcal{S}$ computed with respect to the measure on σ induced by ρ . For $a \in (0, 1]$ define the *intermediate volume growth* of $\sigma \in \mathcal{S}$ by

$$(1) \quad \text{vol}^a(\sigma; \varphi) = \liminf_{n \rightarrow \infty} \frac{\log \text{Vol}(\varphi^n(\sigma))}{n^a} \in [0, \infty],$$

and define the *intermediate volume growth* of φ by

$$\text{vol}^a(\varphi) = \sup_{\sigma \in \mathcal{S}} \text{vol}^a(\sigma; \varphi) \in [0, \infty].$$

Notice that these invariants do not depend on the choice of ρ . Finally define the *volume growth exponent* of φ by

$$\text{exp}_{\text{vol}}(\varphi) := \inf \{a \mid \text{vol}^a(\varphi) < \infty\}.$$

Thus $\text{exp}_{\text{vol}}(\varphi)$ is “the largest $a \in [0, 1]$ such that some submanifold grows under φ like e^{n^a} .” Since $\text{vol}^1(\varphi) \leq (\dim X) \max_{x \in X} \|D\varphi(x)\| < \infty$ we have $\text{exp}_{\text{vol}}(\varphi) \in [0, 1]$. The intermediate volume growth and the volume growth exponent of a smooth flow φ^t on X are defined as $\text{vol}^a(\varphi^1)$ and $\text{exp}_{\text{vol}}(\varphi^1)$.

Remark 1.7. By a celebrated result of Yomdin [35] and Newhouse [22], the volume growth $\text{vol}^1(\varphi)$ agrees with the topological entropy $h_{\text{top}}(\varphi)$. Proceeding as above define for $a \in (0, 1]$ the *intermediate topological entropy* $h_{\text{top}}^a(\varphi)$. We unfortunately do not know whether $\text{vol}^a(\varphi) = h_{\text{top}}^a(\varphi)$ or at least that $\text{vol}^a(\varphi) \leq h_{\text{top}}^a(\varphi)$ also for $a \in (0, 1)$.

Uniform lower bounds for the volume growth or the topological entropy of geodesic flows were found in [5, 24, 25, 27], and these results were generalized to Reeb flows in [19] and, on a polynomial scale, in [8]. Results in [8, 19] show that *the volume growth exponent $\text{exp}_{\text{vol}}(\varphi_\alpha)$ is bounded from below by the maximum of the growth exponent of the function $T \mapsto \text{rank } \iota_*^T H_*(\Omega_0^T M)$ and the growth exponent of the growth function of the fundamental group $\pi_1(M)$* . In particular, for manifolds with fundamental group of exponential growth it is shown in [19] that $\text{vol}^1(\varphi_\alpha) > 0$ for any Reeb flow φ_α on S^*M . Since virtually polycyclic groups are either virtually nilpotent or have exponential growth, we therefore restrict ourselves now to manifolds with virtually nilpotent fundamental group. Since all other fundamental groups of closed manifolds are believed to have exponential growth, this is a minor hypothesis.

Corollary 1.8. *Assume that M is not of finite type, has virtually nilpotent fundamental group, and that there is a field \mathbb{F} such that $H_{\text{m}}(\widetilde{M}; \mathbb{F})$ is infinite-dimensional. Let (S^*M, ξ) be the spherization of M . Then*


$$\text{vol}^{1/2}(S_p^*M; \varphi_\alpha) > 0$$

*for every fiber S_p^*M of S^*M and every Reeb flow φ_α on (S^*M, ξ) . In particular, $\text{vol}^{1/2}(\varphi_\alpha) > 0$ and $\text{exp}_{\text{vol}}(\varphi_\alpha) \geq 1/2$ for every Reeb flow φ_α on (S^*M, ξ) .*

The method. We end this introduction by comparing our approach to previous approaches. As mentioned earlier, lower bounds for the homology of the sublevel sets $\Omega^T M$ follow easily from the growth of the fundamental group $\pi_1(M)$, since its growth is the growth of $H_0(\Omega^T M)$. If $\pi_1(M)$ is finite, Gromov found a way to bound the rank of $H_*(\Omega^T M)$ from below by the rank of $\bigoplus_{i=0}^{cT} H_i(\Omega M)$, where c is a constant depending only on the Riemannian metric. His ingenious argument is purely geometric, see [11, 13, 24], and uses the fact that \widetilde{M} is compact. In [26, 27] Paternain–Petean generalized Gromov’s construction to manifolds with infinite fundamental group, by mapping simply connected complexes K with rich loop space homology into \widetilde{M} . This method gives good lower bounds for the rank of $H_*(\Omega^T M)$ if one can find such complexes and mappings $f: K \rightarrow \widetilde{M}$ for which the homology of $\Omega f(\Omega K)$ still has large rank. This works well for many manifolds, e.g. for most connected sums, for manifolds of finite type with $\pi_*(\Omega M) \otimes \mathbb{Q}$ infinite-dimensional, and in small dimensions, see [26, 27].

To fix the ideas, take as K the wedge sum $S^n \vee S^n$ of two n -spheres. The homology algebra $H_*(\Omega K)$ is the free associative algebra generated by the two generators x_1, x_2 of $H_{n-1}(\Omega K) \cong H_n(K)$, see [4, III. 1. B]. If one can find continuous maps $\iota: K \rightarrow M$ and $r: M \rightarrow K$ such that $r \circ \iota$ is homotopic to the identity of K , then $\Omega \iota: \Omega K \rightarrow \Omega M$ induces an injection in homology. One then concludes that the rank of $\iota_*^T H_*(\Omega^T M)$ grows exponentially. This method was successfully used in dimension ≤ 5 in [27].

For general manifolds of non-finite type, however, we were not even able to prove linear growth of the rank of $H_*(\Omega^T M)$ by this method. Citing G. Paternain, “it is as if one has so much topology that it becomes unmanageable.”



Our strategy to prove lower bounds for the rank of $H_*(\Omega^T M)$ for manifolds of non-finite type is to find an *infinite* wedge sum $\bigvee_{i \in \mathbb{Z}} S_i^n$ of spheres in M as in the figure and to prove that the *ordered* words $x_{i_1} x_{i_2} \cdots x_{i_k}$, $i_1 < i_2 < \cdots < i_k$, are still independent in $H_*(\Omega M; \mathbb{F})$. The proof is neither geometric nor topological, but algebraic. The main point is that we use the Hopf algebra structure of $H_*(\Omega \widetilde{M}; \mathbb{F})$ over the group ring $\mathbb{F}[\pi_1(M)]$, and prove a Poincaré–Birkhoff–Witt type theorem for this algebra. These algebraic results (specifically Theorems 2.1 and 3.6 and Proposition 4.1) are the main findings of this paper. They cover Sections 2 to 4. Theorem 1.1, which is proven in Section 5, follows readily from these algebraic results, and Corollaries 1.5 and 1.8, which are proven in Section 7, follow from Theorem 1.1 by a standard continuation argument in Floer homology and by looking at the volume growth of the Lagrangian disc in $T_p^* M$ bounded by Σ_p .

Acknowledgments. We wish to thank Kenji Fukaya for suggesting to us to bring into play Serre’s Hurewicz theorem, and we thank Andrew Ranicki for his explanations on splitting obstructions. We are grateful to the referees for their suggestions and their careful reading.

The present work is part of the author's activities within CAST, a Research Network Program of the European Science Foundation.

2. HEALTHY G -VECTOR SPACES

Let G be a group, V a vector space over the field \mathbb{F} , and $\rho: G \rightarrow \text{Aut}(V)$ a representation of G . Since the representation is fixed throughout our discussion, we abbreviate $gv = \rho(g)v$ for $g \in G$ and $v \in V$, and we consider the G -vector space V as a module over the group ring $\mathbb{F}G$. A group G is called *polycyclic* if there exists a subnormal series

$$1 = G_0 \triangleleft G_1 \triangleleft \cdots \triangleleft G_n = G$$

such that all factors G_i/G_{i-1} , $1 \leq i \leq n$ are cyclic. A group G is called *virtually polycyclic* if G has a polycyclic subgroup of finite index. Eleven characterisations of polycyclic groups are given in [34, Proposition 4.1]. In this section we prove the following result.

Theorem 2.1. *Assume that G is a virtually polycyclic group acting linearly on an infinite-dimensional vector space V such that V viewed as an $\mathbb{F}G$ -module is finitely generated. Then there exists $v \in V$ and $g \in G$ such that the sequence of vectors $(g^i v)_{i \in \mathbb{Z}}$ is linearly independent.*

Before embarking on the proof of the theorem we introduce some notation. For the following discussion it is irrelevant that G is virtually polycyclic. Given $g \in G$ and $v \in V$ we denote the subvector space of V spanned by the vectors $g^i v$ by

$$W_v^g := \langle g^i v \mid i \in \mathbb{Z} \rangle \subset V.$$

Definition 2.2. A vector $v \in V$ is *healthy* if there exists $g \in G$ such that W_v^g is infinite-dimensional. A vector $v \in V$ which is not healthy is called *sick*.

The reason why we are interested in finding healthy vectors is the following observation from linear algebra.

Lemma 2.3. *If $g \in G$ and $v \in V$, then the following are equivalent.*

- (i) W_v^g is infinite-dimensional.
- (ii) The sequence of vectors $(g^i v)_{i \in \mathbb{Z}}$ is linearly independent.

Proof. That (ii) implies (i) is clear. It remains to show that if (ii) does not hold, then W_v^g is finite-dimensional. The case $v = 0$ is trivial as well, so we assume that $v \neq 0$. Since then $g^{-i}(g^i v) = v \neq 0$, we conclude that $g^i v \neq 0$ for every $i \in \mathbb{Z}$. Hence, if (ii) does not hold, there exist $m < n \in \mathbb{Z}$ and scalars $a_i \in \mathbb{F}$ for $m \leq i \leq n-1$ such that

$$(2) \quad g^n v = \sum_{i=m}^{n-1} a_i g^i v, \quad a_m \neq 0.$$

By applying g^{-m} to (2) we can assume without loss of generality that $m = 0$. Applying g and g^{-1} to (2) we obtain inductively that

$$(3) \quad W_v^g = \langle g^i v \mid 0 \leq i \leq n-1 \rangle.$$

This shows that W_v^g is finite-dimensional. The lemma follows. \square

If $v \neq 0$ is a sick vector, we can define in view of (3) the function $d_v: G \rightarrow \mathbb{N}$ by

$$(4) \quad d_v(g) := \min \left\{ n \in \mathbb{N} \mid W_v^g = \langle g^i v \mid 0 \leq i \leq n-1 \rangle \right\} = \dim W_v^g.$$

Definition 2.4. The G -vector space V is called *sick* if all its vectors are sick. A G -vector space V which is not sick is called *healthy*, i.e., V contains a healthy vector.

Note that a healthy vector space still contains sick vectors. Indeed, the zero vector is always sick. Moreover, observe that the concept of a healthy G -vector space is only of interest if both the cardinality of the group and the dimension of the vector space are infinite, since otherwise V is automatically sick.

Although we fix the representation ρ throughout, we sometimes have to restrict ρ to subgroups $H < G$. In this situation, we say that V is *H -healthy* if the restriction of ρ to H is healthy.

Lemma 2.5. *Assume that $H < G$ is a subgroup of finite index. Then V is G -healthy if and only if V is H -healthy.*

Proof. The implication from H -healthy to G -healthy is obvious. We now assume that V is G -healthy and show that V is H -healthy as well. Since V is G -healthy, there exists a vector $v \in V$ and a group element $g \in G$ such that W_v^g is infinite-dimensional. Denote the right coset gH in G/H by $[g]$.

We first consider the special case where the subgroup H is normal. Then G/H is a group, and its order $n := |G/H|$ is finite by assumption. Hence $[g]^n = id \in G/H$, or equivalently $h := g^n \in H$. By Lemma 2.3 we conclude from the fact that W_v^g is infinite-dimensional that W_v^h is infinite-dimensional as well. This proves that V is H -healthy in the special case that $H \triangleleft G$ is normal.

In the general case where $H < G$ is not necessarily normal, we consider the normal core of H in G defined by

$$\text{Core}(H) := \bigcap_{g \in G} g^{-1}Hg.$$

Note that $\text{Core}(H)$ is a subgroup of H which is normal in G . It is actually the biggest normal subgroup of G contained in H . Moreover, it still has finite index in H , see for instance [29, Theorem 3.3.5]. In view of what we already proved, we therefore conclude that V is $\text{Core}(H)$ -healthy. Since $\text{Core}(H) < H$ it follows that V is H -healthy as well. This finishes the proof of the lemma. \square

If $H < G$ is a subgroup and $v \in V$, we abbreviate by

$$W_v^H = \langle Hv \rangle$$

the subspace of V spanned by the H -orbit of the vector v . The next lemma is our main tool to give an inductive proof of Theorem 2.1.

Lemma 2.6. *Assume that $H \triangleleft G$ is a normal subgroup such that G/H is cyclic. Suppose further that $v \in V$ is sick and W_v^H is finite-dimensional. Then W_v^G is finite-dimensional as well.*

Proof. Abbreviate $m = \dim W_v^H$. Then there exist ξ_1, \dots, ξ_m in the group ring $\mathbb{F}H$ such that

$$(5) \quad W_v^H = \langle \xi_i v \mid 1 \leq i \leq m \rangle.$$

Choose $g \in G$ such that $[g] \in G/H$ is a generator. Assume first that $G/H \cong \mathbb{Z}_n$ is finite. Then

$$W_v^G = \langle g^j \xi_i v \mid 1 \leq i \leq m, 0 \leq j < n \rangle.$$

Hence $\dim W_v^G \leq nm$ is finite. Assume now that $G/H \cong \mathbb{Z}$ is infinite. The case that $v = 0$ is trivial. We therefore assume that $v \neq 0$. Since v is sick by assumption, we have the function $d_v: G \rightarrow \mathbb{N}$ from (4), and we abbreviate $n = d_v(g)$. Our aim is to show that

$$(6) \quad W_v^G = \langle g^j \xi_i v \mid 1 \leq i \leq m, 0 \leq j < n \rangle.$$

For this purpose we abbreviate the right hand side by

$$X := \langle g^j \xi_i v \mid 1 \leq i \leq m, 0 \leq j < n \rangle.$$

That $X \subset W_v^G$ is clear. We have to check the other inclusion, namely that for every $\eta \in \mathbb{F}G$ it holds that

$$\eta v \in X.$$

Since the right coset $[g]$ is a generator of $G/H \cong \mathbb{Z}$, there exist $\eta_j \in \mathbb{F}H$ with $\eta_j \neq 0$ for only finitely many $j \in \mathbb{Z}$ such that

$$\eta = \sum_{j \in \mathbb{Z}} \eta_j g^j.$$

In view of the definition of $n = d_v(g)$, there exists $\zeta \in \mathbb{F}G$ of the form

$$\zeta = \sum_{j=0}^{n-1} \zeta_j g^j, \quad \zeta_j \in \mathbb{F}H$$

such that

$$\eta v = \zeta v.$$

Since H is normal in G , there exist $\zeta'_j \in \mathbb{F}H$ for $0 \leq j < n$ such that

$$\zeta = \sum_{j=0}^{n-1} g^j \zeta'_j.$$

In view of (5) we conclude that for $0 \leq j < n$ we have

$$\zeta'_j v \in \langle \xi_i v \mid 1 \leq i \leq m \rangle.$$

Therefore $\eta v = \zeta v \in X$. This proves (6). We have shown that

$$\dim W_v^G \leq nm = d_v(g) \cdot \dim W_v^H,$$

and therefore W_v^G is finite-dimensional. The proof of the lemma is complete. \square

We are now in position to prove the main result of this section.

Proof of Theorem 2.1. In view of Lemma 2.3 it suffices to show that V is healthy. We argue by contradiction and assume that V is sick. By assumption G is virtually polycyclic, hence contains a polycyclic subgroup H of finite index. By Lemma 2.5 it follows that V is H -sick as well. Since H is polycyclic, we conclude by applying Lemma 2.6 inductively that W_v^H is finite-dimensional for every $v \in V$. Because H has finite index in G it follows that W_v^G is finite-dimensional for every $v \in V$ as well. By assumption V is a finitely generated $\mathbb{F}G$ -module. Therefore we deduce that V is finite-dimensional, contradicting the assumption of the theorem. The proof is complete. \square

3. FILTERED HOPF ALGEBRAS

In this section we consider Hopf algebras V over a group ring $\mathbb{F}G$, endowed with a filtration V^r , $r > 0$, and exhibit a property of V that guarantees that $\dim V^r$ grows at least like $e^{\sqrt{r}}$.

3.1. Hopf algebras over $\mathbb{F}G$. Let G be a group, \mathbb{F} a field and $\mathbb{F}G$ the group ring. We first explain the notion of a Hopf algebra over $\mathbb{F}G$. This might be not completely standard, since usually Hopf algebras are defined over rings which are commutative, a requirement that our group ring in general does not fulfill. However, the feature which distinguishes a group ring from other non-commutative rings is that if V and W are two left modules over $\mathbb{F}G$, then we can still define on their tensor product $V \otimes W = V \otimes_{\mathbb{F}} W$ the structure of a left $\mathbb{F}G$ -module by using the tensor of the two representations: $g(v \otimes w) := (gv) \otimes (gw)$ for $v \in V$, $w \in W$ and $g \in G$. A *product* is then an $\mathbb{F}G$ -linear map $\mu: V \otimes V \rightarrow V$, or equivalently an \mathbb{F} -bilinear map $\mu: V \times V \rightarrow V$ satisfying $\mu(gv, gw) = g\mu(v, w)$ for $g \in G$ and $v, w \in V$. Dually, a *coproduct* is then an $\mathbb{F}G$ -linear map $\Delta: V \rightarrow V \otimes V$. To avoid terrible headaches we assume in addition that our product is always associative, although this requirement is probably not necessary for the results of this section. We abbreviate the product by $vw = \mu(v, w)$. If V in addition is graded, i.e. $V = \bigoplus_{i=0}^{\infty} V_i$, where each V_i is an $\mathbb{F}G$ -submodule of V , then we grade the tensor product by $(V \otimes V)_k = \bigoplus_{i+j=k} V_i \otimes V_j$ and require in addition that the product and coproduct preserve the grading. The product endows the tensor product $V \otimes V$ again with a product which is defined on homogeneous elements by the Koszul sign convention

$$(v \otimes w)(x \otimes y) = (-1)^{\deg(w)\deg(x)} vx \otimes wy$$

where $\deg(v)$ denotes the degree of a homogeneous element v . Given a left module V over $\mathbb{F}G$ and a product μ and coproduct Δ as above, we call the triple (V, μ, Δ) a *bialgebra* over $\mathbb{F}G$ if μ and Δ are compatible in the sense that $\Delta: V \rightarrow V \otimes V$ is a homomorphism of algebras. The bialgebra (V, μ, Δ) is called *connected* if $V_0 = \mathbb{F}$ is one-dimensional and if $1 \in \mathbb{F}$ is also the unit for the multiplication μ .

Definition 3.1. A connected graded bialgebra (V, μ, Δ) over $\mathbb{F}G$ is called a *Hopf algebra* over $\mathbb{F}G$ if for every homogeneous element v of positive degree $\deg(v)$ the coproduct satisfies

$$(7) \quad \Delta v = 1 \otimes v + v \otimes 1 + \sum v_i \otimes v'_i$$

with v_i and v'_i of positive degree. A vector $v \in V$ is *primitive* if $\Delta v = 1 \otimes v + v \otimes 1$.

3.2. Filtrations. In order to endow our Hopf algebra with a filtration we shall suppose that both the vector space V and the group ring $\mathbb{F}G$ are filtered. More precisely, we assume that V can be exhausted by a sequence of finite-dimensional vector spaces, i.e., for every real number $r > 0$ there exists a finite-dimensional subspace $V^r \subset V$ such that $V^r \subset V^s$ for $r \leq s$ and $V = \bigcup_{r>0} V^r$. Define the *value* of $v \in V$ by

$$|v| := \min\{r \mid v \in V^r\}.$$

Notice that for scalars $a_i \in \mathbb{F}$ and vectors $v_i \in V$ we have

$$(8) \quad |a_1 v_1 + \cdots + a_n v_n| \leq \max\{|v_i|\}.$$

Dear reader, please do not confuse the value $|v|$ of v and its degree $\deg(v)$, in case v is homogeneous. They are not related to each other. Also note that the grading is indicated by a subscript while the filtration degree is indicated by a superscript. To get a filtration on the group ring as well, suppose that the group G is endowed with a *length function*, namely a function $L: G \rightarrow \mathbb{R}_{\geq 0}$ satisfying

$$L(g) = L(g^{-1}), \quad L(gh) \leq L(g) + L(h), \quad g, h \in G.$$

In the following we abbreviate $|g| = L(g)$. Via the length function we can define a filtration on the group ring: For $r \geq 0$ we define $\mathbb{F}G^r$ to be the subvector space of $\mathbb{F}G$ consisting of finite sums $\xi = \sum_{g \in G} \xi_g g$ satisfying $\xi_g = 0$ whenever $|g| > r$.

Definition 3.2. The Hopf algebra (V, μ, Δ) over $\mathbb{F}G$ is called *filtered* if the vector space V is endowed with a filtration such that for $v, w \in V$ and $g \in G$ it holds that

$$|vw| \leq |v| + |w|, \quad |gv| \leq |g| + |v|.$$

We next introduce a condition on filtered Hopf algebras which will guarantee nontrivial lower bounds for the growth of $\dim V^r$.

Definition 3.3. Assume that (V, μ, Δ) is a filtered Hopf algebra over $\mathbb{F}G$. A *primitive sequence* is a sequence $(v_i)_{i \in \mathbb{N}}$ satisfying the following conditions.

- (i) The vectors v_i , $i \in \mathbb{N}$, are linearly independent, primitive, and of equal positive degree.
- (ii) There exists a constant $c > 0$ such that $|v_i| \leq ci$.

Remark 3.4. The notion of a primitive sequence does not involve the action of the group G . However, we shall see later that the action of G is useful to construct primitive sequences.

The following result, whose proof is deferred to the next section, will be an important ingredient in the proof of the subsequent Theorem 3.6, which is the main result of this section.

Proposition 3.5. *Let (V, μ, Δ) be a Hopf algebra over \mathbb{F} , and let v_1, \dots, v_N be linearly independent and primitive vectors in V of equal degree. For $I = \{i_1, i_2, \dots, i_\ell\}$ with $1 \leq i_1 < i_2 < \dots < i_\ell \leq N$ write $v_I = v_{i_1} v_{i_2} \cdots v_{i_\ell}$. Then the vectors v_I , $I \subset \{1, \dots, N\}$, are linearly independent.*

Theorem 3.6. *Assume that (V, μ, Δ) is a filtered Hopf algebra over $\mathbb{F}G$ which admits a primitive sequence $(v_i)_{i \in \mathbb{N}}$. Then the function $r \mapsto \dim V^r$ grows at least like $e^{\sqrt{r}}$.*

Proof. Let $I = \{i_1, \dots, i_\ell\} \subset \mathbb{N}$ be a finite subset of distinct numbers which we totally order by $i_1 < i_2 < \dots < i_\ell$, and abbreviate

$$v_I = v_{i_1} v_{i_2} \cdots v_{i_\ell} \in V.$$

Let m be the common degree of the vectors v_i . In view of property (i) of a primitive sequence, Proposition 3.5 shows that the vectors v_I , $I \subset \mathbb{N}$, are linearly independent. We can assume without loss of generality that the constant c in property (ii) of a primitive sequence is 1. Then we have

$$|v_I| \leq \sum_{j=1}^{\ell} |v_{i_j}| \leq \sum_{j=1}^{\ell} i_j.$$

For $k \in \mathbb{N}$ denote by $q(k)$ the number of partitions of k into distinct parts. We have shown that

$$\dim V^n \geq q(1) + q(2) + \cdots + q(n) \geq q(n).$$

By Euler's theorem, the number of partitions of n into distinct parts coincides with the number of its partitions into odd parts, see for example [3, Corollary 1.2]. The asymptotics of this sequence coincides up to a constant with the asymptotics of the partition function (see for example [21, Chapter 16]), which grows like $e^{C\sqrt{n}}$ for a positive constant C according to a theorem of Hardy and Ramanujan, see for example [21, Chapter 15]. \square

Combining Theorems 2.1 and 3.6 we obtain the following result.

Corollary 3.7. *Assume that (V, μ, Δ) is a filtered Hopf algebra over $\mathbb{F}G$ where G is a virtually polycyclic group. Assume further that $\bigoplus_{i < m} V_i$ is finite-dimensional and that V_m is infinite-dimensional but finitely generated as an $\mathbb{F}G$ -module. Then the function $r \mapsto \dim V^r$ grows at least like $e^{\sqrt{r}}$.*

Proof. By Theorem 2.1 there exists $v \in V_m$ and $g \in G$ such that the vectors $v_i := g^i v$, $i \in \mathbb{N}$, are linearly independent. The vectors v_i have equal degree m . Using the defining properties of a filtration, we estimate

$$|v_i| = |g^i v| \leq |g^i| + |v| \leq i|g| + |v|.$$

With $c := |g| + |v|$ we thus have $|v_i| \leq ci$. The sequence $(v_i)_{i \in \mathbb{N}}$ therefore meets all the properties of a primitive sequence, except that the v_i may fail to be primitive. To correct this, we consider the linear map $A: \langle v_1, v_2, \dots \rangle \rightarrow V_m \otimes V_m$ given by

$$A(v) = \Delta v - 1 \otimes v - v \otimes 1.$$

Since the vector space $\bigoplus_{j < m} V_j$ is finite-dimensional, the subvector space of $V_m \otimes V_m$ spanned by the elements $u_i \otimes u'_i$ with $u_i, u'_i \in \bigoplus_{j < m} V_j$ is finite-dimensional. Since A takes values in this finite-dimensional vector space, $k := \text{rank } A$, which equals the dimension of $\langle v_1, v_2, \dots \rangle / \ker A$, is finite. In particular, $\ker A$ is infinite-dimensional. We shall construct by induction a sequence w_1, w_2, \dots of linearly independent elements in $\ker A$ such that $|w_i| \leq c(k+1)$. The sequence $(w_i)_{i \in \mathbb{N}}$ is then a primitive sequence in V , and the corollary follows in view of Theorem 3.6.

The restriction of the map A to $\langle v_1, v_2, \dots, v_{k+1} \rangle$ has a non-trivial kernel. Let $w_1 := a_1 v_1 + \dots + a_{k+1} v_{k+1}$ be a non-trivial element in this kernel. Then by (8), $|w_1| \leq c(k+1)$. Next, the restriction of A to $\langle v_{k+2}, v_{k+3}, \dots, v_{2(k+1)} \rangle$ has a non-trivial kernel. Let $w_2 := a_{k+2} v_{k+2} + \dots + a_{2(k+1)} v_{2(k+1)}$ be a non-trivial element in this kernel. Then by (8), $|w_2| \leq 2c(k+1)$, and w_1, w_2 are linearly independent because $v_1, \dots, v_{2(k+1)}$ are linearly independent. Proceeding in this way we construct linearly independent vectors w_1, w_2, \dots such that $|w_i| \leq ic(k+1)$. \square

4. A QUANTUM POINCARÉ–BIRKHOFF–WITT THEOREM

Consider a graded bialgebra (V, μ, Δ) over the field \mathbb{F} which is connected (i.e., $V_0 = \mathbb{F}$ is one-dimensional) and is such that $1 \in \mathbb{F}$ also serves as the unit for the multiplication μ . (The group G plays no role in this section.) We again assume that μ is associative, and write $vw = \mu(v, w)$. Also recall that $v \in V$ is *primitive* if $\Delta(v) = 1 \otimes v + v \otimes 1$. Suppose that for $N \in \mathbb{N}$ we are given linearly independent and primitive elements $v_1, \dots, v_N \in V_m$ of equal positive degree $m \in \mathbb{N}$. Abbreviate $\mathbb{N}_N = \{1, \dots, N\}$. We order $I \subset \mathbb{N}_N$ using the canonical order of \mathbb{N}_N , namely we write $I = \{i_1, \dots, i_\ell\}$ satisfying $i_1 < i_2 < \dots < i_\ell$. We then abbreviate $v_I = v_{i_1} v_{i_2} \dots v_{i_\ell}$, where we use the convention that $v_\emptyset = 1$. The following proposition reminiscent of the Poincaré–Birkhoff–Witt theorem seems to be known to people working in quantum group theory, cf. [31, Theorem 1.5(b)]. However, since the Hopf algebras arising in the theory of quantum groups are usually not graded, we provide a proof for the readers convenience.

Proposition 4.1. *Assume that v_1, \dots, v_N are linearly independent and primitive vectors in V . Then the vectors v_I , $I \subset \mathbb{N}_N$, are linearly independent.*

Proof. The crucial ingredient in the proof is the computation of the coproduct of the elements v_I . To determine the signs in this formula, we use the following convention. Given a subset I of $\{1, \dots, N\}$ we order $I = \{i_1, \dots, i_\ell\}$ and its complement $I^c = \{j_1, \dots, j_{N-\ell}\}$. This determines a permutation $(1, \dots, N) \mapsto (i_1, \dots, i_\ell, j_1, \dots, j_{N-\ell})$. We denote by $\sigma(I)$ the signum of this permutation.

Lemma 4.2. *The coproduct of $v_{\mathbb{N}_N}$ is given by*

$$\Delta(v_{\mathbb{N}_N}) = \sum_{I \subset \mathbb{N}_N} \sigma(I)^m v_I \otimes v_{I^c}.$$

Hence if m is even, then $\Delta(v_{\mathbb{N}_N}) = \sum_{I \subset \mathbb{N}_N} v_I \otimes v_{I^c}$, and if m is odd, then $\Delta(v_{\mathbb{N}_N}) = \sum_{I \subset \mathbb{N}_N} \sigma(I) v_I \otimes v_{I^c}$.

Proof. We prove the lemma by induction on N . For $N = 1$ the lemma is an immediate consequence of the fact that v_1 is primitive. For the induction step we assume that the formula holds for $N - 1$. We compute

$$\begin{aligned} \Delta(v_{\mathbb{N}_N}) &= \Delta(v_{\mathbb{N}_{N-1}} v_N) \\ &= \Delta(v_{\mathbb{N}_{N-1}}) \Delta(v_N) \\ &= \left(\sum_{I \subset \mathbb{N}_{N-1}} \sigma(I)^m v_I \otimes v_{\mathbb{N}_{N-1} \setminus I} \right) \left(1 \otimes v_N + v_N \otimes 1 \right) \\ &= \sum_{I \subset \mathbb{N}_{N-1}} \sigma(I)^m v_I \otimes (v_{\mathbb{N}_{N-1} \setminus I} v_N) \\ &\quad + \sum_{I \subset \mathbb{N}_{N-1}} (-1)^{\deg(v_{\mathbb{N}_{N-1} \setminus I}) \deg(v_N)} \sigma(I)^m (v_I v_N) \otimes v_{\mathbb{N}_{N-1} \setminus I}. \end{aligned}$$

Since $\deg(v_N) = m$ and $\deg(v_{\mathbb{N}_{N-1} \setminus I}) = m(N - 1 - \#I)$, we have for each $I \subset \mathbb{N}_{N-1}$ that

$$\begin{aligned} (-1)^{\deg(v_{\mathbb{N}_{N-1} \setminus I}) \deg(v_N)} \sigma(I)^m &= (-1)^{m^2(N-1-\#I)} \sigma(I)^m \\ &= (-1)^{m(N-1-\#I)} \sigma(I)^m \\ &= \sigma(I \cup \{N\})^m. \end{aligned}$$

The previous sum therefore becomes

$$\begin{aligned} &= \sum_{I \subset \mathbb{N}_{N-1}} \sigma(I)^m v_I \otimes v_{\mathbb{N}_N \setminus I} \\ &\quad + \sum_{I \subset \mathbb{N}_{N-1}} \sigma(I \cup \{N\})^m v_{I \cup \{N\}} \otimes v_{\mathbb{N}_N \setminus (I \cup \{N\})} \\ &= \sum_{I \subset \mathbb{N}_N} \sigma(I)^m v_I \otimes v_{\mathbb{N}_N \setminus I}. \end{aligned}$$

This proves the induction step and hence the lemma. \square

Proof of Proposition 4.1. Recall that by assumption the vectors v_1, \dots, v_N are linearly independent and have all the same degree m . Since $\deg v_I = m^{\#I}$ it suffices to show that for every $k \leq N$ the vectors v_I with $I \subset \mathbb{N}_N$ and $\#I = k$ are linearly independent. For $k = 1$ this is an assumption. For the induction step we assume this assertion for all $k \leq n - 1$ where $n \leq N$. Since the coproduct is linear, it suffices to show that the vectors $\Delta(v_I)$ with $I \subset \mathbb{N}_N$, $\#I = n$ are linearly independent. It follows from the induction

hypothesis that for every $0 < k < n$ the vectors $v_I \otimes v_J$ with $I, J \subset \mathbb{N}_N$ and $\#I = k$, $\#J = n - k$ are linearly independent. This and the formulas

$$\Delta(v_I) = \sum_{J \subset I} \sigma(J)^m v_J \otimes v_{I \setminus J},$$

that follow from (the proof of) Lemma 4.2, complete the induction step. \square

5. PROOF OF THEOREM 1.1

We shall deduce Theorem 1.1 from Corollary 3.7. We start with saying briefly “who is who” in Corollary 3.7: Let M be a closed connected manifold, and let \widetilde{M} be its universal covering space. We take as G the fundamental group $\pi_1(M)$ of M , and as V we take the homology $H_*(\Omega\widetilde{M}; \mathbb{F})$ over a suitable field \mathbb{F} . The product μ will be the Pontryagin product given by concatenation of loops, and the coproduct will simply come from the diagonal map $\Omega\widetilde{M} \rightarrow \Omega\widetilde{M} \times \Omega\widetilde{M}$, $x \mapsto (x, x)$. Now fix a Riemannian metric on M . The filtration on V^r will be given by taking $|v|$ as the smallest r for which v can be represented by a cycle of based loops of length at most r , and for $g \in \pi_1(M)$ we take $L(g)$ to be (half of) the length of the shortest curve representing g .

We shall show in Sections 5.1 and 5.2 that with these choices, (V, μ, Δ) is a filtered Hopf algebra over $\mathbb{F}G$. In Section 5.3 we show that for M of non-finite type with $G = \pi_1(M)$ virtually polycyclic, the assumption in Theorem 1.1 that $H_m(\widetilde{M}; \mathbb{F})$ is infinite-dimensional implies the dimension assumptions on $\oplus V_i$ in Corollary 3.7.

5.1. The Hopf-algebra structure on $H_*(\Omega\widetilde{M}; \mathbb{F})$. Let M be a closed connected manifold, and let $\text{pr}: \widetilde{M} \rightarrow M$ be its universal covering space. Fix $p \in M$ and $\tilde{p} \in \widetilde{M}$ over p . The spaces $\Omega_0 M$ and $\Omega\widetilde{M}$ of contractible continuous loops based at p and \tilde{p} , respectively, are canonically identified. Conjugation of loops in $\Omega_0 M$ by loops in M based at p yields an action of the fundamental group $G = \pi_1(M, p)$ on $H_*(\Omega_0 M) = H_*(\Omega\widetilde{M})$: Given a cycle $C = \{\gamma\}$ of loops in $\Omega\widetilde{M}$ based at \tilde{p} and given $g \in G$, the class $g[C]$ is defined as the class represented by the cycle of loops $\{c_g^{-1} \circ g\gamma \circ c_g\}$, where c_g is the lift to \widetilde{M} starting at \tilde{p} of a loop in M in class g , and $g\gamma$ is the lift of $\text{pr} \circ \gamma$ starting at $g\tilde{p}$.

Let \mathbb{F} be the field, and abbreviate $V_* := H_*(\Omega\widetilde{M}; \mathbb{F})$. The action of G on V extends to an action of $\mathbb{F}G$ on V . Concatenation of loops in \widetilde{M} based at \tilde{p} induces a product $\mu: V \otimes V \rightarrow V$, called the Pontryagin product. Since concatenation of loops is associative up to homotopy, μ is associative. It follows from the definition of the action of G that μ is $\mathbb{F}G$ -linear. In order to define the coproduct $\Delta: V \rightarrow V \otimes V$, we consider, more generally, a topological space X and the diagonal map $\delta_X: X \rightarrow X \times X$, $x \mapsto (x, x)$. Since we work over a field \mathbb{F} , the cross product

$$H_*(X; \mathbb{F}) \otimes H_*(X; \mathbb{F}) \xrightarrow{\times} H_*(X \times X; \mathbb{F})$$

is an isomorphism by the Künneth formula. We can therefore define $\Delta_X: H_*(X; \mathbb{F}) \rightarrow H_*(X; \mathbb{F}) \otimes H_*(X; \mathbb{F})$ by

$$\Delta_X := \times^{-1} \circ (\delta_X)_*.$$

Assume now in addition that X is a path-connected H-space, with product ν .

Lemma 5.1. *The homology $H_*(X; \mathbb{F})$, with product induced by ν and with coproduct Δ_X , is a Hopf algebra.*

We refer to [33, Theorem 7.15] for the proof. For the readers convenience, we verify that for every homogeneous element v of positive degree, $\Delta_X v$ has the form (7). Let $p: X \times X \rightarrow X$, $(x, y) \mapsto x$, be the projection on the first factor. Then

$$(9) \quad p \circ \delta_X = id_X.$$

For an element $u = v'_n \otimes 1 + 1 \otimes v''_n + \sum v'_i \otimes v''_j \in \bigoplus_{i+j=n} H_i(X) \otimes H_j(X)$ with $\deg v'_i < n$ we have

$$(10) \quad (p_n \circ \times)u = v'_n$$

by the geometric definition of the cross product (see e.g. [15, §3.B]). Now write $\Delta_X v = v'_n \otimes 1 + 1 \otimes v''_n + \sum v'_i \otimes v''_j$ with $\deg v'_i, \deg v''_j \geq 1$. Using (9), the definition of Δ_X and (10) we get

$$\begin{aligned} v &= p_n \circ (\delta_X)_n v \\ &= p_n \circ \times \circ \Delta_X v \\ &= p_n \circ \times \left(v'_n \otimes 1 + 1 \otimes v''_n + \sum v'_i \otimes v''_j \right) \\ &= v'_n. \end{aligned}$$

Similarly we find $v''_n = v$, and so $\Delta_X v = v \otimes 1 + 1 \otimes v + \sum v'_i \otimes v''_j$ with $\deg v'_i, \deg v''_j \geq 1$. \square

Since \widetilde{M} is simply connected, $\Omega\widetilde{M}$ is path-connected. Hence $V_0 = H_0(\Omega\widetilde{M}; \mathbb{F}) \cong \mathbb{F}$ is one-dimensional. Moreover, $1 \in \mathbb{F}$ corresponds to the class of the constant path $\tilde{p} \in \widetilde{M}$, which is the unit for the Pontryagin product μ . Applying Lemma 5.1 with $X = \Omega\widetilde{M}$ and writing Δ for $\Delta_{\Omega\widetilde{M}}$ we obtain that (V, μ, Δ) is a Hopf algebra over $\mathbb{F}G$. (We have already noticed that μ is $\mathbb{F}G$ -linear. The definition of the G -actions on V and on $V \otimes V$ shows that also Δ is $\mathbb{F}G$ -linear.)

5.2. The filtration on $H_*(\Omega\widetilde{M}; \mathbb{F})$. Fix a Riemannian metric ρ on M , and let $\tilde{\rho} = \text{pr}^* \rho$ be the corresponding Riemannian metric on \widetilde{M} . Given a piecewise smooth curve γ in \widetilde{M} , we denote by $\ell(\gamma)$ the length of γ with respect to the Riemannian metric $\tilde{\rho}$. For $r > 0$ let V^r be the set of homology classes in $V = H_*(\Omega\widetilde{M}; \mathbb{F})$ that can be represented by cycles formed by piecewise smooth loops γ based at \tilde{p} with $\ell(\gamma) \leq r$,

$$V^r := \iota_*^r H_*(\Omega^r \widetilde{M}; \mathbb{F}).$$

Then each V^r is finite-dimensional (see [20, §16]), $V^r \subset V^s$ for $r \leq s$ and $V = \bigcup_{r>0} V^r$. As in Section 3.2 define the value of $v \in V$ by $|v| := \min \{r \mid v \in V^r\}$. In view of the definition of the Pontryagin product and by the triangle inequality, $|vw| \leq |v| + |w|$ for all $v, w \in V$.

Next, for $g \in G = \pi_1(M, p)$ let $\ell(g)$ be the minimal length of a piecewise smooth loop based at p that represents g . In other words, $\ell(g)$ is the length of the shortest geodesic lasso based at p in class g . Set $L(g) := \frac{1}{2}\ell(g)$. Then $L(g) = L(g^{-1})$ and $L(gh) \leq L(g) + L(h)$ for all $g, h \in G$ by the triangle inequality. Finally, for $g \in G$ denote by \underline{c}_g the lift to \widetilde{M} based at \tilde{p} of a shortest curve in class g . Then

$$\ell(\underline{c}_g^{-1} \circ g\gamma \circ \underline{c}_g) \leq \ell(\underline{c}_g^{-1}) + \ell(g\gamma) + \ell(\underline{c}_g) = \ell(\gamma) + 2\ell(\underline{c}_g) = \ell(\gamma) + L(g)$$

for all $g \in G$ and $\gamma \in \Omega\widetilde{M}$. In view of the definition of the G -action on $\Omega\widetilde{M}$ we find that $|gv| \leq |v| + L(g) = |v| + |g|$. We have shown that (V, μ, Δ) is a filtered Hopf algebra over $\mathbb{F}G$.

5.3. Dimensions. Recall that M is of *finite type* if its universal cover \widetilde{M} is homotopy equivalent to a finite CW-complex.

Lemma 5.2. ([8, Lemma 2.2]) *The following are equivalent.*

- (i) M is of finite type.
- (ii) The Abelian groups $H_k(\widetilde{M})$ are finitely generated for all $k \geq 1$.
- (iii) The Abelian groups $\pi_k(M)$ are finitely generated for all $k \geq 2$.

Now assume that M is not of finite type. By the lemma, we can define

$$(11) \quad \mathfrak{m}(M) := \min \left\{ k \mid H_k(\widetilde{M}) \text{ is not finitely generated} \right\} \in \{2, \dots, \dim M\}.$$

The main result of this subsection is

Proposition 5.3. *Assume that M is not of finite type. Let $\mathfrak{m} = \mathfrak{m}(M)$ be as in definition (11). Then*

- (i) $H_i(\Omega\widetilde{M})$ is finitely generated for $i \leq \mathfrak{m} - 2$;
- (ii) $H_{\mathfrak{m}-1}(\Omega\widetilde{M})$ is not finitely generated, but is finitely generated as a $\mathbb{Z}G$ -module.

Proof. For each $k \geq 2$ the fundamental group $G = \pi_1(M, p)$ acts on $\pi_k(M) = \pi_k(M, p)$ by conjugation. Under the Hurewicz homomorphism $h: \pi_k(M) = \pi_k(\widetilde{M}) \rightarrow H_k(\widetilde{M})$ this action corresponds to the action of G on $H_k(\widetilde{M})$ induced by deck transformations (the commutative diagram on the left). On $\pi_{k-1}(\Omega\widetilde{M}) = \pi_{k-1}(\Omega_0 M) \cong \pi_k(M)$ this action of G is induced by conjugation of elements in $\Omega_0 M \cong \Omega\widetilde{M}$. This action also induces an action on $H_{k-1}(\Omega\widetilde{M})$, namely the action described in Section 5.1, and the two actions commute with the Hurewicz homomorphism $h: \pi_{k-1}(\Omega\widetilde{M}) \rightarrow H_{k-1}(\Omega\widetilde{M})$ (the commutative diagram

on the right),

$$\begin{array}{ccccc}
H_k(\widetilde{M}) & \xleftarrow{h} & \pi_k(\widetilde{M}) = \pi_{k-1}(\Omega\widetilde{M}) & \xrightarrow{h} & H_{k-1}(\Omega\widetilde{M}) \\
\downarrow & & \downarrow & & \downarrow \\
H_k(\widetilde{M}) & \xleftarrow{h} & \pi_k(\widetilde{M}) = \pi_{k-1}(\Omega\widetilde{M}) & \xrightarrow{h} & H_{k-1}(\Omega\widetilde{M})
\end{array}$$

Recall that $\mathfrak{m} \in \{2, \dots, \dim M\}$ is the minimal integer such that $H_{\mathfrak{m}}(\widetilde{M})$ is not finitely generated. By Serre's theory of \mathcal{C} -classes, applied to the class of finitely generated Abelian groups, \mathfrak{m} is also the minimal integer such that $\pi_{\mathfrak{m}}(\widetilde{M})$ is not finitely generated. More precisely, Serre's Hurewicz theorem implies that for $k \leq \mathfrak{m}$ the Hurewicz map $h: \pi_k(\widetilde{M}) \rightarrow H_k(\widetilde{M})$ is injective and surjective up to finitely generated groups, see [30] or [16, Theorem 1.8]. Hence $\pi_k(\Omega\widetilde{M})$ is finitely generated for $k \leq \mathfrak{m} - 2$, but not so for $k = \mathfrak{m} - 1$.

Since $\Omega\widetilde{M}$ is a path-connected H-space, $\pi_1(\Omega\widetilde{M})$ acts trivially on $\pi_k(\Omega\widetilde{M})$ for $k \geq 0$. Serre's Hurewicz theorem now implies that $h: \pi_k(\Omega\widetilde{M}) \rightarrow H_k(\Omega\widetilde{M})$ has finitely generated kernel and cokernel for $k \leq \mathfrak{m} - 1$, see [30, p. 274] or [16, Theorem 1.8]. It follows that $H_k(\Omega\widetilde{M})$ is finitely generated for $k \leq \mathfrak{m} - 2$, but not so for $k = \mathfrak{m} - 1$.

We are left with proving the second assertion in (ii). After replacing M by a homotopy equivalent space, if necessary, we find a CW-structure on M . Since M is compact, this CW-structure is finite. We lift this structure to \widetilde{M} by the action of G . The cellular chain complex $C_*(\widetilde{M}; \mathbb{Z})$ is then a finitely generated $\mathbb{Z}G$ -module in each degree. Since G is virtually polycyclic, the ring $\mathbb{Z}G$ is left Noetherian, see [14] or [17]. Hence each $\mathbb{Z}G$ -module $C_*(\widetilde{M}; \mathbb{Z})$ is left Noetherian. Therefore the kernel and the image of the differential of $C_*(\widetilde{M}; \mathbb{Z})$ as well as the quotient $H_*(\widetilde{M})$ are finitely generated left Noetherian $\mathbb{Z}G$ -modules in each degree. In particular, $H_{\mathfrak{m}}(\widetilde{M})$ is a finitely generated $\mathbb{Z}G$ -module. Recall that $h: \pi_{\mathfrak{m}}(\widetilde{M}) \rightarrow H_{\mathfrak{m}}(\widetilde{M})$ is injective and surjective up to finitely generated groups. In view of the commutative diagram above, it follows that $\pi_{\mathfrak{m}}(\widetilde{M})$ and hence $\pi_{\mathfrak{m}-1}(\Omega\widetilde{M})$ are finitely generated $\mathbb{Z}G$ -modules. As we have seen before, $h: \pi_{\mathfrak{m}-1}(\Omega\widetilde{M}) \rightarrow H_{\mathfrak{m}-1}(\Omega\widetilde{M})$ has finitely generated cokernel. Hence $H_{\mathfrak{m}-1}(\Omega\widetilde{M})$ is also a finitely generated $\mathbb{Z}G$ -module. \square

5.4. Proof of Theorem 1.1 and Remark 1.2.1. By the universal coefficient theorem,

$$(12) \quad H_i(\Omega\widetilde{M}; \mathbb{F}) = H_i(\Omega\widetilde{M}) \otimes \mathbb{F} \oplus \text{Tor}(H_{i-1}(\Omega\widetilde{M}); \mathbb{F})$$

for every field \mathbb{F} . Hence assertion (i) of Proposition 5.3 implies that $H_i(\Omega\widetilde{M}; \mathbb{F})$ is finite-dimensional for $i \leq \mathfrak{m} - 2$, and assertion (ii) implies that $H_{\mathfrak{m}-1}(\Omega\widetilde{M}; \mathbb{F})$ is finitely generated as an $\mathbb{F}G$ -module. Assume now that \mathbb{F} is a field such that $H_{\mathfrak{m}}(\widetilde{M}; \mathbb{F})$ is infinite-dimensional. Since

$$(13) \quad H_{\mathfrak{m}}(\widetilde{M}; \mathbb{F}) = H_{\mathfrak{m}}(\widetilde{M}) \otimes \mathbb{F} \oplus \text{Tor}(H_{\mathfrak{m}-1}(\widetilde{M}); \mathbb{F})$$

and since $H_{m-1}(\widetilde{M})$ is finitely generated, $H_m(\widetilde{M}) \otimes \mathbb{F}$ is also infinite-dimensional. As we have seen in the proof above, the two Hurewicz maps

$$(14) \quad H_m(\widetilde{M}) \xleftarrow{h} \pi_m(\widetilde{M}) = \pi_{m-1}(\Omega\widetilde{M}) \xrightarrow{h} H_{m-1}(\Omega\widetilde{M})$$

both have finitely generated kernel and cokernel. It follows that $H_{m-1}(\Omega\widetilde{M}) \otimes \mathbb{F}$ is also infinite-dimensional. This and (12) imply that $H_{m-1}(\Omega\widetilde{M}; \mathbb{F})$ is infinite-dimensional. Hence the dimension assumptions in Corollary 3.7 are satisfied, and we conclude that $\dim V^T$ grows at least like $e^{\sqrt{T}}$.

Similarly, (13) shows that $\dim H_m(\widetilde{M}; \mathbb{F}) = \infty$ if and only if $\dim H_m(\widetilde{M}) \otimes \mathbb{F} = \infty$, which holds if and only if $\dim \pi_m(\widetilde{M}) \otimes \mathbb{F} = \infty$ because the left map in (14) has finitely generated kernel and cokernel. \square

6. A GENERALIZATION TO SIMPLICIAL COMPLEXES, AND A ‘‘COUNTEREXAMPLE’’

6.1. An extension to simplicial complexes. Our main result Theorem 1.1 holds true for CW-complexes. To avoid some technicalities with rectifiable loops, we restrict ourselves to simplicial complexes, linearly embedded in some Euclidean space.

Consider a finite simplicial complex K . Fix a piecewise linear embedding of K into some \mathbb{R}^N . Then every point $x \in K$ has a neighbourhood $U \subset K$ such that for every $u \in U$ the straight line from x to u lies in K and is strictly shorter than any other rectifiable curve in K from x to u . Milnor’s method of approximating the loop space by broken geodesics, [20, §16], now shows that every cycle in ΩK is homotopic to a cycle in the space of piecewise linear loops in K . Hence every homology class in the loop space ΩK can be represented by a cycle formed by piecewise linear loops.

Denote by $\Omega_0^T K$ the space of contractible based loops in K that are piecewise linear and of length $\ell(\gamma) \leq T$. For further use we state two other consequences of the approximation method in [20, §16] (see also [13, p. 353]): First, each sublevel $\Omega_0^T K$ is homotopy equivalent to a finite polyhedron, and hence

$$(15) \quad \iota_*^T H_*(\Omega_0^T K) \quad \text{is finitely generated for each } T.$$

In particular, all vector spaces $\iota_*^T H_*(\Omega_0^T K; \mathbb{F})$ are finite-dimensional. Second, there exists a constant $c = c(X, \rho)$ such that no k -cycle in $\Omega_0 K$ can be represented by a cycle in $\Omega_0^{ck} K$. In other words, for every $k \geq 1$,

$$(16) \quad \iota_k^T H_k(\Omega_0^T K) = 0 \quad \text{for } T \leq ck.$$

The proof of Theorem 1.1 applies literally to the present situation:

Theorem 6.1. *Let K be a finite connected simplicial complex, embedded into some \mathbb{R}^N . Assume that K is not of finite type and has virtually polycyclic fundamental group. Assume further that there exists a field \mathbb{F} such that $H_m(\widetilde{K}; \mathbb{F})$ is infinite-dimensional. Then the function of T*

$$\dim \iota_*^T H_*(\Omega_0^T K; \mathbb{F})$$

grows at least like $e^{\sqrt{T}}$.

6.2. A “counterexample”. We now illustrate the role of the assumption that there exists a field \mathbb{F} such that $H_{\mathfrak{m}}(\tilde{K}; \mathbb{F})$ is infinite-dimensional by an example. Let X be the mapping torus of a degree- p map f of the sphere S^n , where p is prime and $n \geq 2$. Since self-maps of spheres of equal degree are homotopic, p and n determine X up to homotopy equivalence. If we endow S^n with the CW-structure with two cells, the 0-cell \star being a fixed point of f , then the attaching map f is cellular, and we obtain a CW-structure on X . Since $H_k(X) = \mathbb{Z}$ for $k = 0, 1$ and $H_k(X) = 0$ otherwise, X is not homotopy-equivalent to a topological manifold.

We now compute $H_*(\tilde{X})$ and $H_*(\Omega\tilde{X})$. The fundamental group $\pi_1(X) = \mathbb{Z}$ is cyclic. The universal cover \tilde{X} is the “double mapping telescope” obtained by glueing together the mapping cylinders of $f_i = f$, $i \in \mathbb{Z}$. For integers $i \leq j$ consider the part of \tilde{X} over $[i, j]$,

$$\tilde{X}_{[i,j]} = \{(x, t) \in \tilde{X} \mid t \in [i, j]\},$$

and write $\tilde{X}_j = \tilde{X}_{[j,j]}$. By moving along t we see that $\tilde{X}_{[i,j]}$ deformation retracts onto $\tilde{X}_j = S^n$. Retract $\tilde{X}_{[-1,0]}$ to \tilde{X}_0 in time $[0, \frac{1}{2}]$. Then $\tilde{X}_{[-2,-1]} \subset \tilde{X}$ becomes $\tilde{X}_{[-1,0]}$; retract it to \tilde{X}_0 in time $[\frac{1}{2}, \frac{3}{4}]$. Going on this way we construct a deformation retraction $\tilde{X} \rightarrow \tilde{X}_{\geq 0}$ onto the mapping telescope formed by the mapping cylinders of $f_i = f$, $i \geq 0$.

Since f has degree p , the inclusion $\tilde{X}_{[0,j]} \rightarrow \tilde{X}_{[0,j+1]}$ induces multiplication by p in homology,

$$H_n(\tilde{X}_{[0,j]}) = \mathbb{Z} \xrightarrow{p} \mathbb{Z} = H_n(\tilde{X}_{[0,j+1]}).$$

The direct limes of the sequence $\mathbb{Z} \xrightarrow{p} \mathbb{Z} \xrightarrow{p} \mathbb{Z} \rightarrow \cdots$ can be identified with $\mathbb{Z}[1/p]$, the subgroup of \mathbb{Q} consisting of rational numbers with denominators a power of p . Hence

$$H_n(\tilde{X}) = H_n(\tilde{X}_{\geq 0}) = \varinjlim H_n(\tilde{X}_{[0,j]}) \cong \mathbb{Z}[1/p],$$

(cf. Example 3.F.3 in [15]), and $H_k(\tilde{X}) = 0$ for all other $k \geq 1$. The Abelian group $\mathbb{Z}[1/p]$ is not finitely generated, hence $\mathfrak{m} = n$ in this example. However, for a field \mathbb{F} of characteristic q we have $H_n(\tilde{X}; \mathbb{F}) = 0$ if $q = p$ and $H_n(\tilde{X}; \mathbb{F}) \cong \mathbb{F}$ otherwise.

We next compute $H_*(\Omega\tilde{X})$. We choose as base point of \tilde{X} the point $\star \in \tilde{X}_0 \cong S^n$. Let \mathfrak{p} be the path from $\star \in \tilde{X}_0$ to $(\star, j) \in \tilde{X}_j$ traced by \star under the canonical deformation retraction $\tilde{X}_{[0,j]} \rightarrow \tilde{X}_j$. The space $(\Omega\tilde{X}_{[0,j]}, \star)$ is homeomorphic to the subspace of $(\Omega\tilde{X}_{[0,j]}, (\star, j))$ obtained by conjugating by \mathfrak{p} the loops in $(\Omega\tilde{X}_{[0,j]}, (\star, j))$. Under the deformation retraction $\tilde{X}_{[0,j]} \rightarrow \tilde{X}_j$ this subspace deformation retracts onto $(\Omega\tilde{X}_j, (\star, j)) = (\Omega S^n, \star)$. We have

$$H_k(\Omega S^n) = \begin{cases} \mathbb{Z} & \text{if } k \in (n-1)\mathbb{N} \cup 0, \\ 0 & \text{otherwise.} \end{cases}$$

Fix $k = (n-1)\ell$. The natural homotopy equivalence $J(S^{n-1}) \rightarrow \Omega S^n$ from the James reduced product of S^{n-1} to ΩS^n shows that the map $\Omega f: \Omega S^n \rightarrow \Omega S^n$ induces multiplication

by p^ℓ in H_k ,

$$H_k(\Omega S^n) = \mathbb{Z} \xrightarrow{p^\ell} \mathbb{Z} = H_k(\Omega S^n),$$

see [18] or [15, §4.J]. The direct limes of the sequence $\mathbb{Z} \xrightarrow{p^\ell} \mathbb{Z} \xrightarrow{p^\ell} \mathbb{Z} \longrightarrow \dots$ can be identified with $\mathbb{Z}[1/p^\ell]$, the subgroup of \mathbb{Q} consisting of rational numbers with denominators a power of p^ℓ . Hence

$$H_k(\Omega \tilde{X}) = H_k(\Omega \tilde{X}_{\geq 0}) = \varinjlim H_k(\Omega \tilde{X}_{[0,j]}) \cong \mathbb{Z}[1/p^\ell].$$

Moreover, $H_k(\Omega \tilde{X}) = 0$ if $k \notin (n-1)\mathbb{N} \cup 0$.

Every finite CW-complex is homotopy equivalent to a finite simplicial complex, see [32, Theorem 24]. So choose a finite simplicial complex K homotopy equivalent to X , and fix an embedding of K into some Euclidean space.

Proposition 6.2. *The space K fulfills all assumptions of Theorem 6.1 with $\mathfrak{m} = n$, besides for $H_{\mathfrak{m}}(\tilde{K}; \mathbb{F})$ being zero- or one-dimensional for every field \mathbb{F} . Moreover, $\iota_*^T H_*(\Omega_0^T K)$ is a free Abelian group whose rank grows not more than linearly with T .*

Proof. Since X and K are homotopy equivalent, so are \tilde{X} and \tilde{K} . The homologies of \tilde{X} and \tilde{K} are thus isomorphic, and so are the homologies of $\Omega \tilde{X}$ and $\Omega \tilde{K}$. It remains to understand the growth of $\iota_*^T H_*(\Omega^T \tilde{K})$. If $k \notin (n-1)\mathbb{N} \cup 0$, then $\iota_k^T H_k(\Omega^T \tilde{K}) \subset H_k(\Omega \tilde{K}) = 0$, so $\iota_k^T H_k(\Omega^T \tilde{K}) = 0$. If $k = (n-1)\ell$, then $\iota_k^T H_k(\Omega^T \tilde{K}) \subset H_k(\Omega \tilde{K}) = \mathbb{Z}[1/p^\ell]$. By (15), the group $\iota_k^T H_k(\Omega^T \tilde{K})$ is finitely generated. Every finitely generated subgroup of $\mathbb{Z}[1/p^\ell]$ is contained in the subgroup generated by some element $1/p^{i\ell}$ and hence is trivial or infinite cyclic. Moreover, by (16) there exists a constant c depending only on K and ρ such that $\iota_k^T H_k(\Omega^T \tilde{K}) = 0$ for $T \leq ck$. Hence $\iota_*^T H_*(\Omega^T \tilde{K})$ is a free Abelian group whose rank grows not more than linearly with T . \square

7. PROOF OF COROLLARIES 1.5 AND 1.8

Proof of Corollary 1.5. Let \mathcal{C} be a component of $\Omega_{pq} M$. Fix a smooth path $c \in \mathcal{C}$, of length $\ell(c)$. The map $h: \Omega_0(M, p) \rightarrow \mathcal{C}$, $\gamma \mapsto c \circ \gamma$ is a homotopy equivalence. It maps $\Omega_0^T M$ to $\mathcal{C}^{T+\ell(c)}$, where \mathcal{C}^T is the space of piecewise smooth paths in \mathcal{C} of length $\leq T$. Theorem 1.1 now implies that

$$(17) \quad \dim \iota_*^T H_*(\mathcal{C}^T; \mathbb{F}) \text{ grows at least like } e^{\sqrt{T}}.$$

Notice that $\dim \iota_*^T H_*(\mathcal{C}^T; \mathbb{F}) \leq \dim H_*(\mathcal{C}^T; \mathbb{F})$. For geodesic flows, Corollary 1.5 now follows from classical Morse theory, see [20, Theorem 16.3] or [25, p. 116]. For the general case of Reeb flows, we use (17) and Lagrangian Floer homology, exactly as in [19, Section 6]. We outline the argument, omitting all technical issues.

Let $\Sigma \subset T^*M$ be the fiberwise starshaped hypersurface corresponding to the cooriented contact manifold (S^*M, α) , and (up to the time change $t \mapsto 2t$) view the Reeb flow φ_α^t on (S^*M, α) as the restriction of the Hamiltonian flow φ_K^t on T^*M , where $\Sigma = K^{-1}(1)$ and K is fiberwise homogeneous of degree 2.

Denote by $\mathcal{P}(K, \mathcal{C})$ the set of smooth paths $x: [0, 1] \rightarrow T^*M$ with $x(0) \in \Sigma_p$, $x(1) \in \Sigma_q$ that are solutions to Hamilton's equation $\dot{x}(t) = X_K(x(t))$ and whose projections to M belong to \mathcal{C} . The action of $x \in \mathcal{P}(K, \mathcal{C})$ is

$$\mathcal{A}_K(x) = \int_0^1 (\lambda(\dot{x}(t)) - K(x(t))) dt,$$

where λ is the Liouville 1-form on T^*M . Since K is homogenous of degree two, the set $\mathcal{P}^{T^2/2}(K, \mathcal{C}) = \{x \in \mathcal{P}(K, \mathcal{C}) \mid \mathcal{A}_K(x) \leq T^2/2\}$ is in bijection with the Reeb chords from Σ_p to Σ_q of time $\leq T$. We thus wish to show that $\#\mathcal{P}^{T^2/2}(K, \mathcal{C})$ grows at least like $e^{\sqrt{T}}$.

Set $a = T^2/2$. Since p, q are non-conjugate for the Reeb flow φ_α^t , the set $\mathcal{P}^a(K, \mathcal{C})$ is finite. The Floer chain group $\text{CF}^a(K, \mathcal{C})$ is the \mathbb{F} -vector space freely generated by the chords in $\mathcal{P}^a(K, \mathcal{C})$. The Conley–Zehnder index of these chords (normalized such that it agrees with the Morse index in case of a non-degenerate geodesic chord) gives this vector space a grading $*$. The Floer boundary operator on $\text{CF}_*^a(K, \mathcal{C})$ is a map of degree -1 . Its homology is the Floer homology $\text{HF}_*^a(K, \mathcal{C})$. Since the dimension of the homology $\text{HF}_*^a(K, \mathcal{C})$ is not greater than the dimension of the chain group $\text{CF}_*^a(K, \mathcal{C})$, which equals $\#\mathcal{P}^a(K, \mathcal{C})$, we wish to show that $\dim \text{HF}_*^a(K, \mathcal{C})$ grows at least like $e^{\sqrt{T}}$.

Fix a Riemannian metric g on M , and let $G: T^*M \rightarrow \mathbb{R}$ be the corresponding geodesic Hamiltonian. After multiplying g with a constant, we can assume that $G \leq K$, and since Σ is fiberwise starshaped with respect to the origin, we find a constant $\sigma \geq 1$ such that

$$G \leq K \leq \sigma G.$$

The Floer homology of the functions G and σG is defined in the same way.

There is a commutative diagram

$$\begin{array}{ccccc} \text{HF}_*^a(G, \mathcal{C}) & \xrightarrow{\Phi_{G \sigma G}} & \text{HF}_*^{a/\sigma}(\sigma G, \mathcal{C}) & \xrightarrow{\text{ASM}} & H_*(\mathcal{C}^T; \mathbb{F}) \\ \Phi_{GH} \downarrow & & \downarrow & & \downarrow \\ \text{HF}_*^a(K, \mathcal{C}) & \xrightarrow{\Phi_{K \sigma G}} & \text{HF}_*^a(\sigma G, \mathcal{C}) & \xrightarrow{\text{ASM}} & H_*(\mathcal{C}^{\sqrt{\sigma}T}; \mathbb{F}) \end{array}$$

Here, the three maps Φ between Floer homologies are Floer continuation maps, and $\Phi_{G \sigma G}$ is an isomorphism. The upper map ASM is the composition

$$\text{HF}_*^{a/\sigma}(\sigma G, \mathcal{C}) \xrightarrow{\text{AS}} \text{HM}_*^{a/\sigma}(L, \mathcal{C}) \xrightarrow{\text{AM}} H_*(\mathcal{C}^T; \mathbb{F})$$

of the Abbondandolo–Schwarz isomorphism from Floer homology to the Morse homology of the Legendre transform L of σG with the Abbondandolo–Majer isomorphism from this homology to the homology of \mathcal{C}^T , see [1] and [2]. Finally, the two unlabeled vertical arrows are induced by inclusion.

It follows that $\dim \text{HF}_*^a(K, \mathcal{C})$ is at least the rank of the right vertical map. This rank is at least $\dim \iota_*^T H_*(\mathcal{C}^T; \mathbb{F})$, which by (17) grows at least like $e^{\sqrt{T}}$. \square

Proof of Corollary 1.8. For geodesic flows the claim follows from Corollary 1.5 and the geometric arguments in [25, Section 3.1]. For the general case of Reeb flows we use Theorem 1.1 and the idea from [6], that was further developed in [7, 8, 19]. The proof goes exactly as the proof of Theorem 4.6 in [19]; we therefore again only sketch the proof.

For $q \in M$ consider $\Sigma_q = \Sigma \cap T_q^*M$ and the bounded component D_q of $T_q^*M \setminus \Sigma_q$. The “spheres” Σ_q are Legendrian and the “discs” D_q are Lagrangian. Fix $n \in \mathbb{N}$, and take $q \in M$ not conjugate to p for the flow φ_α^t . Recall from the previous proof that the generators of the Floer chain groups $\text{CF}_*^{n^2/2}(K)$ (where we now do not restrict to a component \mathcal{C}) are in bijection with the Reeb chords from Σ_p to Σ_q of time $\leq n$. This set is in bijection with the intersection points of the Lagrangian discs $\varphi_K^n(D_p)$ and D_q . By the previous proof, this number grows at least like $e^{\sqrt{n}}$, uniformly in q . Since this holds for almost every $q \in M$, the volume of the discs $\varphi_K^n(D_p)$ grows at least like $e^{\sqrt{n}}$. We refer to Section 4 of [19] for details. Finally, the volume of $\varphi_K^n(\Sigma_p)$ also grows like $e^{\sqrt{n}}$ in view of Proposition 4.3 in [8]. \square

REFERENCES

- [1] A. Abbondandolo and P. Majer. Lectures on the Morse complex for infinite-dimensional manifolds. Morse theoretic methods in nonlinear analysis and in symplectic topology, 1–74, *NATO Sci. Ser. II Math. Phys. Chem.*, **217**, Springer, Dordrecht, 2006.
- [2] A. Abbondandolo and M. Schwarz. On the Floer homology of cotangent bundles. *Comm. Pure Appl. Math.* **59** (2006) 254–316.
- [3] G. Andrews. *The theory of partitions*, Cambridge University Press (1976).
- [4] R. Bott and H. Samelson. On the Pontryagin product in spaces of paths. *Comment. Math. Helv.* **27** (1953), 320–337 (1954).
- [5] E. I. Dinaburg. A connection between various entropy characterizations of dynamical systems. *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971) 324–366.
- [6] U. Frauenfelder and F. Schlenk. Volume growth in the component of the Dehn–Seidel twist. *Geom. Funct. Anal.* **15** (2005) 809–838.
- [7] U. Frauenfelder and F. Schlenk. Fiberwise volume growth via Lagrangian intersections. *J. Symplectic Geom.* **4** (2006) 117–148.
- [8] U. Frauenfelder, C. Labrousse and F. Schlenk. Slow volume growth for Reeb flows on spherizations and contact Bott–Samelson theorems. arXiv:1307.7290
- [9] R. Grigorchuk. Groups of intermediate growth: an introduction. *Enseign. Math.* **54** (2008) 251–272.
- [10] R. Grigorchuk. On the Gap Conjecture concerning group growth. arXiv:1202.6044
- [11] M. Gromov. Homotopical effects of dilatation. *J. Differential Geom.* **13** (1978) 303–310.
- [12] M. Gromov. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.* **53** (1981), 53–73.
- [13] M. Gromov. *Metric structures for Riemannian and non-Riemannian spaces*. Modern Birkhäuser Classics. Birkhäuser Boston, Boston, 2007.
- [14] P. Hall. Finiteness conditions for soluble groups. *Proc. London Math. Soc.* **4** (1954) 419–436.
- [15] A. Hatcher. Algebraic topology. *Cambridge University Press, Cambridge*, 2002.
- [16] A. Hatcher. Spectral Sequences in Algebraic Topology. www.math.cornell.edu/~hatcher
- [17] S. V. Ivanov. Group rings of Noetherian groups. *Mat. Zametki* **46** (1989), no. 6, 61–66, 127; translation in *Math. Notes* **46** (1989), no. 5–6, 929–933 (1990).
- [18] I. M. James. Reduced product spaces. *Ann. of Math.* **62** (1955) 170–197.

- [19] L. Macarini and F. Schlenk. Positive topological entropy of Reeb flows on spherizations. *Math. Proc. Cambridge Philos. Soc.* **151** (2011) 103–128.
- [20] J. Milnor. Morse theory. *Annals of Mathematics Studies* **51**, Princeton University Press, Princeton, 1963.
- [21] M. Nathanson. *Elementary methods in number theory*, Graduate Texts in Mathematics **195** (2000).
- [22] S. Newhouse. Entropy and volume. *Ergodic Theory Dynam. Systems* **8*** (1988), Charles Conley Memorial Issue, 283–299.
- [23] V. Ozols. Largest normal neighborhoods. *Proc. Amer. Math. Soc.* **61** (1976) 99–101 (1977).
- [24] G. Paternain. Topological entropy for geodesic flows on fibre bundles over rationally hyperbolic manifolds. *Proc. Amer. Math. Soc.* **125** (1997) 2759–2765.
- [25] G. Paternain. *Geodesic flows*. Progress in Mathematics **180**. Birkhäuser Boston, Inc., Boston, MA, 1999.
- [26] G. Paternain and J. Petean. Entropy and collapsing of compact complex surfaces. *Proc. London Math. Soc.* **89** (2004) 763–786.
- [27] G. Paternain and J. Petean. Zero entropy and bounded topology. *Comment. Math. Helv.* **81** (2006) 287–304.
- [28] A. Ranicki. Exact sequences in the algebraic theory of surgery. *Mathematical Notes* **26**. Princeton University Press, Princeton; University of Tokyo Press, Tokyo, 1981.
- [29] W. Scott. *Group theory*, Prentice-Hall, Englewood Cliffs, New Jersey (1964).
- [30] J.-P. Serre. Groupes d’homotopie et classes de groupes abéliens. *Ann. of Math.* **58** (1953) 258–294.
- [31] D. Wang, J. Zhang, and G. Zhuang. Lower bounds of growth of Hopf algebras, *Trans. Amer. Math. Soc.* **365**, 4963–4986 (2013).
- [32] J. H. C. Whitehead. A certain exact sequence. *Ann. of Math.* **52** (1950) 51–110.
- [33] G. W. Whitehead. *Elements of homotopy theory*. Graduate Texts in Mathematics **61**. Springer-Verlag, New York-Berlin, 1978.
- [34] J. Wolf. Growth of finitely generated solvable groups and curvature of Riemannian manifolds. *J. Differential Geometry* **2** (1968) 421–446.
- [35] Y. Yomdin. Volume growth and entropy. *Israel J. Math.* **57** (1987) 285–300.

URS FRAUENFELDER, DEPARTMENT OF MATHEMATICS AND RESEARCH INSTITUTE OF MATHEMATICS,
SEOUL NATIONAL UNIVERSITY
E-mail address: frauenf@snu.ac.kr

FELIX SCHLENK, INSTITUT DE MATHÉMATIQUES, UNIVERSITÉ DE NEUCHÂTEL
E-mail address: schlenk@unine.ch